



THE UNIVERSITY *of* EDINBURGH

## Edinburgh Research Explorer

### Categorical speech perception in adults with autism spectrum conditions

**Citation for published version:**

Stewart, ME, Petrou, AM & Ota, M 2018, 'Categorical speech perception in adults with autism spectrum conditions', *Journal of Autism and Developmental Disorders*, vol. 48, no. 1, pp. 72-82.  
<https://doi.org/10.1007/s10803-017-3284-0>

**Digital Object Identifier (DOI):**

[10.1007/s10803-017-3284-0](https://doi.org/10.1007/s10803-017-3284-0)

**Link:**

[Link to publication record in Edinburgh Research Explorer](#)

**Document Version:**

Peer reviewed version

**Published In:**

Journal of Autism and Developmental Disorders

**Publisher Rights Statement:**

The final publication is available at Springer via <http://dx.doi.org/10.1007/s10803-017-3284-0>

**General rights**

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

**Take down policy**

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact [openaccess@ed.ac.uk](mailto:openaccess@ed.ac.uk) providing details, and we will remove access to the work immediately and investigate your claim.



## **Categorical Speech Perception in Adults with Autism Spectrum Conditions**

Mary E. Stewart <sup>1</sup>, Alexandra M. Petrou <sup>1,2</sup>, Mitsuhiro Ota <sup>3</sup>

<sup>1</sup> Heriot-Watt University, Edinburgh, Scotland, EH14 4AS

<sup>2</sup> Institute of Neuroscience, Newcastle University, Newcastle Upon Tyne, England, NE1 4LP

<sup>3</sup> Philosophy, Psychology and Language Sciences, University of Edinburgh, Edinburgh,  
Scotland, EH8 9AD

## Abstract

This study tested whether individuals with autism spectrum conditions (n=23) show enhanced discrimination of acoustic differences that signal a linguistic contrast (i.e., /g/ versus /k/ as in ‘goat’ and ‘coat’) and whether they process such differences in a less categorical fashion as compared with 23 IQ-matched typically developed adults. Tasks administered were nonverbal IQ, verbal IQ, 5 language measures, a speech perception task, and the ADOS. The speech perception task measured the discrimination of paired exemplars along the /g/-/k/ continuum. Individuals with autism spectrum conditions did not show enhanced discrimination of speech perception. Categorical speech perception was correlated with verbal ability of reading, lexical decision, and verbal IQ in individuals with ASC.

Key Words: Categorical speech perception; autism; auditory discrimination; language; phoneme.

Individuals with Autism Spectrum Conditions (ASC) exhibit perceptual processing patterns that systematically differ from those of typically developed individuals. On the one hand, individuals with ASC demonstrate enhanced perceptual performance in both auditory and visual domains, including pitch processing (Bonnell et al. 2003; Heaton 2003; Heaton 2005; Heaton et al. 1998; Heaton et al. 1999; Mottron et al. 2000; Järvinen-Pasley et al. 2008; Stewart et al. 2015) and visual pattern recognition (Plaisted et al. 1998). On the other hand, they show deficits in higher-order cognitive tasks that involve the same perceptual domains in which they show superior performance, such as the interpretation of intonational phrases or detection of emotion from faces and speech (Ashwin et al. 2007; Boraston et al. 2007; Korpilahti et al. 2007; Kujala et al. 2005; McCann and Peppé 2003). These paradoxical findings have been interpreted as manifestations of the manner in which individuals with ASC are thought to coordinate different levels of processing, namely, low levels that deal directly with the physical properties of the stimuli, and high levels where the percept of the stimuli are linked to relevant mental representations (e.g., intonational meanings, facial expressions). For instance, the style of processing associated with ASC may show: weakened integration of lower level processing with higher level processing (i.e., Weak Central Coherence; Frith 2003); an increased level of lower level processing (i.e., Enhanced Perceptual Functioning; Mottron and Burack 2001); or a lack of flexibility in ignoring prediction errors, which leads to a tendency to focus on local processing at the expense of more abstract representations of the stimuli (Van de Cruys et al. 2014).

It has been proposed that such diminished interaction between levels of processing should result in less categorical perception of stimuli (Soulières et al. 2007). Categorical perception occurs when percepts that vary continuously along a physical dimension are perceived as more

1 distinct when they are mapped onto separate than onto the same category. A consequence of  
2 categorical perception is that a stimulus pair that spans the boundary between two categories is  
3 more discriminable than one that belongs to the same category. If the cognitive style associated  
4 with ASC means that there is a reduced role of cognitive categories in the processing of the  
5 physical properties of the stimuli, this boundary effect should be attenuated in individuals with  
6 ASC. Indeed, this is what Soulières et al. (2007) found for visual shape perception. Participants  
7 in their study were presented with tokens of ellipsis that had a constant height but varying  
8 widths. In a same-different task, participants without ASC showed heightened discriminability  
9 of stimulus pairs drawn from the midpoint region of the ellipsis continuum, the presumed  
10 location of the boundary for the two categories of ellipses, whereas individuals with ASC did  
11 not exhibit a comparable peak in discrimination around that region.  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28

29 If this finding from Soulières et al. (2007) is a reflection of a broadly diminished influence of  
30 categories on perception in ASC, we expect to find similar effects in other processing domains  
31 such as speech sounds, where categorical perception has been robustly demonstrated in  
32 typically developing individuals (Liberman et al. 1957; Liberman et al. 1967). Categorical  
33 perception of speech sounds offers an important extension of the finding from ellipsis  
34 discrimination by Soulières et al. (2007) for two reasons. Firstly, humans do not normally have  
35 pre-established categories for ellipses, so the boundary-like effect observed in Soulières et al.'s  
36 participants without ASC must have come from emergent categories that they acquired through  
37 exposure to the stimulus set in the experiment. Stimulus repetition contributes to reduce  
38 discrimination of within-category differences whilst enhance between-category discrimination  
39 (Harnad 1987). Therefore one interpretation of the Soulières et al. (2007) study is that there is  
40 slower category formation in individuals with ASC. In contrast, speech sounds, such as /g/ and  
41 /k/ in *goat* and *coat*, tend to form categories that are firmly in place in adult speakers. By  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

testing speech discrimination, we can therefore examine whether ASC affects categorical perception of pre-existing contrasts.

Secondly, if observed, diminished top-down processing in the perception of speech sounds provides one possible explanation for some of the language-related problems reported for individuals with ASC (Kjelgaard and Tager-Flusberg 2001). For a listener who has continuous or noncategorical perception of the phonetic dimension that distinguishes, say, /g/ and /k/, the between-category sound difference that separates words such as *goat* and *coat* would be less distinct. The same listener may also be riveted by within-category acoustic differences that are not relevant to such lexical contrasts; an effect that is likely to interfere with his or her linguistic processing. Reduced categorical perception offers a testable link between auditory perception skills and language functions in individuals with ASC, which to this date, has not been fully established (although see Jones et al. 2009 and Järvinen-Pasley et al. 2008).

In the current study, we examined the effects of ASC on categorical speech perception by comparing individuals with and without ASC on their discrimination of auditory stimuli taken from a continuum exemplifying the contrast between the segment /g/ and /k/. The main acoustic cue to a voicing contrast that can be heard in syllables such as /gɪ/-/kɪ/ is the temporal lag between the release burst of the consonant and the onset of the periodic wave of the following vowel: the voice-onset time (VOT). Hypothetical discrimination curves of VOT are illustrated in Figure 1. A typical discrimination curve for stimulus pairs on a VOT continuum has a peak in sensitivity around the boundary between the two sound categories - this is what would be expected from the typically developed sample and illustrated by line (a) in Figure 1. If categorical discrimination is reduced, a discrimination curve such as line (b) is hypothesised. Methodologically, the difference between these two curves can be captured by comparing the

1 difference in discrimination level between the centre of the continuum (30/50 ms in Figure. 1)  
2 and the endpoints of the continuum (e.g, 10/30 and 50/70 ms in Figure. 1). This measure,  
3  
4 which we call the *Categorical Discrimination Index*, should be smaller in individuals with  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
Individuals with ASC may have a generally high level of discrimination ability, which causes  
their discrimination curve to look ‘flatter’ due to a ceiling effect. This scenario is illustrated by  
line (c) in Figure 1. To fully understand the characteristics of speech perception in ASC, we  
therefore need to examine both the general level of sensitivity (i.e., mean discrimination level)  
and the shape of sensitivity (i.e., the Categorical Discrimination Index). The main objective of  
our study was to use these measures to compare individuals with and without ASC on their  
discrimination curves for a /gɪ/-/kɪ/ VOT continuum.

(Figure 1 here)

The second objective of our study was to examine whether characteristics of speech perception  
in ASC are related to aspects of language functioning. As discussed above, less categorical  
perception of speech sounds may result in less clear-cut processing of the phonetic information  
signaling phonemic differences such as /g/ versus /k/. This may adversely affect phonological  
working memory capacity (e.g., remembering novel words such as *glate*, as opposed to *clate*)  
as well as identification of known words (e.g., recognizing *goat*, as opposed to *coat*). Deficits  
in phonological representations, in turn, have been associated with problems in orthographic  
learning and processing (Swan and Goswami 1997; Snowling 2000). Following these lines of

reasoning, we investigated whether the degree to which speech sound perception is categorical is related to measures of nonword repetition, vocabulary knowledge, reading and spelling.

One caveat in studying perceptual patterns in relation to ASC is that performance in discrimination tasks can be confounded with IQ. Although a recent study has reported that perceptual ability in individuals with ASC shows no correlation with IQ (Meilleur et al. 2014), relationships between discrimination performance and IQ have been observed in typically developed adults (Deary 1994). In order to control for potential confounding effects of IQ, we recruited only adults with ASC who had been identified as being average or above in terms of their IQ and typically developing individuals who were individually matched on nonverbal-IQ.

## Methods

### Participants

Twenty-three adults with ASC (mean age = 25.8 years, SD = 7.2, range = 18–40 years; males = 15, females = 8) were compared with 23 nonverbal-IQ matched typically-developed adults (mean age = 28.7 years, SD = 9.8, range = 18–49 years; males = 12, females = 11). Participants were included based on a clinical diagnosis established prior to recruitment according to DSM-IV-TR criteria (APA, 2000) through multidisciplinary assessment by clinical services. In order to further characterise the participants, participants with ASC also completed the Autism Diagnostic Observation Schedule (ADOS; Lord et al. 2000) and all participants completed the Autism-Spectrum Quotient (AQ; Baron-Cohen et al. 2001) as part of the current study. Participants were classified into the ASC group if they had a diagnosis and if they met the cut-off scores on the ADOS (Lord et al. 2000) or the AQ (Woodbury-Smith et al. 2005).



Each adult with ASC was matched on a one-to-one basis to a typically-developed adult exactly on their nonverbal-IQ score using the Raven's Advanced Progressive Matrices (meaning that the participants had exactly the same nonverbal IQ score, the participants will be referred to as IQ-matched adults throughout; Raven et al. 2003). Participants were also matched as closely as possible on gender. A strategy of controlling for IQ was taken, instead of the alternative strategy of recruiting a sample of adults who varied in IQ and later controlling for IQ, due to the nature of the task and the difficulties of testing a similar paradigm across a wide IQ range.

The adults with ASC were recruited from universities and from drop-in and support centres for individuals with ASC. There was no significant difference in the number of males or females between the adults with ASC and the IQ matched adults ( $\chi^2[1] = 0.81, p = .37$ ). All participants had English as their primary native language and a score above 4 on the Raven's Advanced Progressive Matrices. A score of 4 was chosen as a cut-off point because participants with a lower score might not have fully engaged with the test. None of the participants met the following exclusion criteria for the studies: history of neurological disorder, genetic or medical problems that might affect hearing, any known learning disability including speech and language impairments, and individuals who were unable to provide informed consent.

## Materials

### *Tests of speech perception*

An ABX discrimination task and an auditory lexical decision task were adapted from Stewart and Ota (2008). The ABX discrimination task was constructed using a 5-step continuum that was created using the Klatt synthesiser (Sensyn version 1.1). We created 7 base stimuli that varied in VOT from 10 ms to 70 ms in 10 ms steps and made to sound like utterances ranging

from /gI/ to /kI/. All base stimuli had a duration of 250 ms and an identical rise-fall fundamental frequency contour. The VOT was manipulated by incrementally increasing the amplitude of voicing at the end of the aspiration interval from 0 to 64 dB, the amplitude of aspiration from 0 to 60 dB and the bandwidth of the first formant from 60 to 400 Hz for the duration of the aspiration interval. We then created the ABX stimuli by concatenating two base stimuli 20 ms apart (e.g., 10 and 30 ms), with an inter-stimulus interval of 1 second. The resulting ABX stimuli had VOT steps of 10-30 ms, 20-40 ms, 30-50 ms, 40-60 ms, and 50-70 ms.

The task comprised two blocks of trials. Each block consisted of 4 randomised permutations (i.e., ABA, ABB, BBA, BAB) of the 8 sets of ABX stimuli. Thus, each VOT step was tested 8 times (i.e., 4 times x 2 blocks). The practice session comprised 6 trials. Participants were required to listen to the recorded stimuli and press the numbers '1' or '2' on the keyboard to indicate whether the third sample was the same as the first sample or the second sample, respectively. Test time took approximately 10 minutes. Accuracy and reaction time were recorded.

We also asked whether adults with ASC would be less categorical in their discrimination of basic speech sounds. This would be observed as a less pronounced discrimination peak at the boundary in the ASC group compared to IQ-matched adults (see Figure 1 for hypothetical discrimination curves). This hypothesis was tested by an index of 'categoricalness', or Categorical Discrimination Index (CD Index), which we calculated by taking the mean accuracy of responses of the two endpoint stimuli (i.e., the mean of Step 10-30 ms and Step 50-70 ms) and subtracting this value from the accuracy of response at the presumed boundary in the continuum, where typically developed individuals characteristically demonstrate peak

performance (i.e., boundary Step 30-50 ms). A lower CD Index indicates a smaller disparity between between-category discrimination and within-category discrimination and, thus, reduced categorical speech perception.

### *Auditory lexical decision*

Stimuli were 48 monosyllable real words and nonwords which were read by a male speaker. The critical experimental items were pairs of words that differ by the voicing of the initial consonant such that one member was a real word in English (e.g., *golf*) and the other was not (e.g., *colf*). The place of articulation ([*p/b*], [*t/d*], and [*k/g*]) and voice/voiceless direction were balanced. Half of the tokens were fillers, half of which were real words and half nonwords. Each item was played once in a random order to comprise 48 unblocked trials. Participants were required to listen to the recorded stimuli and press the ‘y’ or ‘n’ keys on the keyboard to indicate whether the item they heard was a real word (“Yes”) or not a real word (“No”), respectively.

### *Nonverbal IQ*

Test set II of Raven’s Advanced Progressive Matrices (Raven et al. 2003), which consists of 36 items, was used. Each item is multiple choice, with eight options. Answering an item correctly gives a score of 1, and incorrectly a score of 0. Therefore, the range of possible scores is 0–36. The task items were presented by a computer using the E-prime package (Psychological Software Tools, Pittsburgh, PA). We shortened the test in order to reduce participant fatigue. Rather than select a subset of items, to preserve the progressive nature of the task, we chose to present the full task with a 20 minute time limit (see Fugard et al. 2011). A high correlation is shown between this method and the full test score ( $r = .74$ ; Hamel and Schmittmann 2006).

### *Verbal IQ*

The Millhill Vocabulary Scales (Raven et al. 2003), which measure acquired verbal knowledge, were used. The test comprises two lists of words split into two sets of 34 words with the first item treated as a practice item within each set. Administered as a pen-and-paper task, the participant is required to select the synonymous word from a list of 6 possible choices. The difficulty of items increases throughout the test. A correct answer gives a score of 1 and an incorrect answer gives a score of 0, with a possible score range of 0–33.

### *Reading and Spelling*

The Reading and Spelling components of the Wide Range Achievement Test – Third Revision (WRAT-3; Wilkinson 1993) were used. For the Reading task, participants are required to recognise and pronounce 42 words out of context. If no response is given within 10 seconds, a prompt is given to continue to the next word on the list. The Word Spelling task comprises 40 words. Each word is required to be spelled to dictation. In both cases, a correct answer gives a score of 1 and an incorrect answer gives a score of 0.

### *Phonological short-term memory*

Phonological short-term memory was measured using a test developed from previously designed nonword repetition tasks (e.g. Grube et al. 2013). A list of nonwords comprising 20 items ranging from 4 to 8 syllables in length was played to the participants. There were 4 practice items, all of which were 4 syllables in length. The participant was required to repeat the nonword after hearing it. Two nonword repetition scores were computed for each participant: A total correct score and a nonword repetition accuracy score. The total correct score was calculated by taking the total number of correctly repeated nonwords out of 20. The

nonword repetition accuracy score was calculated by taking the mean percentage of correct sounds repeated across all 20 nonwords.

### *Autism-Spectrum Quotient*

The Autism-Spectrum Quotient (AQ; Baron-Cohen et al. 2001) is made up of 50 questions, which are divided into 5 theoretically defined subscales: *social skill*, *attention switching*, *attention to detail*, *communication*, and *imagination*. Each item has four possible answers: ‘strongly agree’, ‘agree’, ‘disagree’, and ‘strongly disagree’. We used Likert scoring (e.g., Austin 2005; Clark et al. 2013; Hoekstra et al. 2007; Stewart and Ota 2008) and included all four levels when computing scores. Item scores ranged from 1–4, so total AQ scores ranged from 50–200, and the subscores ranged from 10–40, with a higher score indicating more autistic traits.

### *Autism Diagnostic Observation Schedule*

The Autism Diagnostic Observation Schedule (ADOS; Lord et al. 2000) is a standardised assessment of symptoms associated with the autism spectrum for autism diagnosis and classification. In this study, module 4, which is appropriate for adults who are verbally fluent, was administered. Scores are derived for diagnostic algorithms of Communication, Social Interaction, and Restricted/Stereotyped Interests, Behaviours or Activities.

### *Procedure*

Participants were given the experimental tasks in the following order: ABX nonword discrimination, auditory lexical decision, AQ, Raven’s, Millhill, Reading, Spelling, and nonword repetition. There was missing data for 4 adults with ASC and 4 typically-developed adults for the nonword repetition task: one participant with ASC did not wish to be audio

1 recorded which was necessary to carefully listen back and score nonwords; for the remaining 7  
2 participants there were technical difficulties with the audio recording which meant it could not  
3 be scored. The ADOS was administered to the adults with ASC following this battery or on a  
4 separate occasion. If a separate occasion was required, this occurred no longer than 1 week  
5 subsequent to the first session. Adults were individually tested in a quiet lab or at an  
6 appropriate place at their institution and received payment for their participation. Overall test  
7 time took approximately 2 hours and 10 mins for the adults with ASC and 1 hour and 20 mins  
8 for the typically-developed adults.  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21

22 The ABX nonword discrimination, auditory lexical decision, and Raven's tasks were delivered  
23 using the stimulus presentation programme E-Prime (Psychological Software Tools, Inc.).  
24

25 Stimuli were played on a Toshiba, Satellite Pro laptop over headphones (Edirol Audio Capture  
26 UA-3FX; Sennheiser HD 265). All the speech sounds and lexical decision task was adjusted to  
27 a comfortable volume for the individual.  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38

### 39 Data preparation

40  
41 To ensure that responses to items on each of the tasks were from those trials to which  
42 participants attended, items were removed where RTs were greater than 2.5 SD above the  
43 participant's mean latency for each auditory task. For adults with ASC, this resulted in the  
44 removal of 30 items (2.3%) from the Discrimination Task and 33 items (3.0%) from the  
45 Lexical Decision Task. For IQ-matched adults, this resulted in the removal of 51 items (3.7%)  
46 from the Discrimination Task and 35 items (2.9%) from the Lexical Decision Task.  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

## Results

Table 1 displays the descriptive statistics of the AQ Likert Total, the Raven's, the Millhill, the number of participants who completed each task, and the language measures for the adults with ASC and the IQ-matched adults. Adults with ASC scored significantly higher on the AQ Likert Total than IQ-matched adults ( $t[44] = 5.02, p = .000$ , Cohen's  $d = 1.5$ ). There was no significant difference between the groups on the Raven's, as the participants were exactly matched on this measure. There were no significant group differences on any of the other measures. One participant, who was identified as having a diagnosis of ASC, did not meet the ADOS or the AQ cut-off scores. The analyses were repeated with and without this matched pair. Inclusion or exclusion of this matched pair did not affect the significance of the results throughout except in reference to the comparison of the Categorical Discrimination Index, analyses with and without the matched pair are included only in this instance, and are outlined below. All other analyses do not include this participant and their matched control.

(Table 1 here)

#### *Is discrimination of basic speech sounds enhanced in adults with ASC?*

We tested whether individuals with ASC would show a generally higher level of VOT discrimination compared to IQ matched controls (see Figure 2). There was no significant main effect of Group ( $F[1, 44] = 0.065, p = .80$ ), or Group x Step interaction ( $F[4, 176] = 1.75, p = .14$ ). Response time was also assessed in order to test whether the groups approached the task in a different way (see Figure 3). There was no significant main effect of Group ( $F[1, 44] = 0.03, p = .85$ ), or Group by Step interaction ( $F[4, 176] = 0.68, p = .61$ ). Both groups therefore appear to process the stimuli in a similar way. This analysis did not provide support for the idea that the general level of discrimination of basic speech sounds is enhanced in adults with ASC.

(Figure 2 here)

(Figure 3 here)

### *Is discrimination of basic speech sounds less categorical in adults with ASC?*

An independent-samples t-test revealed that adults with ASC had a lower CD Index than the IQ-matched adults. This was statistically significant with the additional matched pair ( $t[46] = 2.07, p = .04, M = 0.09, SD = 0.31$  for adults with ASC;  $M = 0.24, SD = 0.17$  for IQ-matched adults, Cohen's  $d = 0.60$ ) and a similar effect size and a trend were found in an analysis without the additional pair ( $t[44] = 1.94, p = .06, M = 0.10, SD = 0.31$  for adults with ASC;  $M = 0.25, SD = 0.18$  for IQ-matched adults, Cohen's  $d = 0.59$ ). A similar index was calculated for response time. There was no significant difference between the groups ( $t[44] = 1.42, p = .16$ ). To verify the results of the CD Index analysis of the discrimination accuracy, we also compared the kurtoses of the accuracy scores between the two groups. The distribution of accuracy of discrimination across the 5 VOT Steps was more platykurtic for adults with ASC (Kurtosis =  $-0.11, SE = 0.94$ ) than IQ-matched adults (Kurtosis =  $-1.19, SE = 0.94; t[44] = 2.36, p = .02$ ). Therefore, these analyses provide moderate support for the idea that adults with ASC are less categorical in their discrimination of basic speech sounds.

### *Correlations with language measures and with symptom severity*

We tested whether the CD Index correlated with performance on the language measures of Reading, Spelling, Lexical Discrimination, Vocabulary, and phonological awareness (see Table 2) in adults with ASC and IQ-matched adults. The CD Index correlated significantly with Reading, Lexical Discrimination and Vocabulary in individuals with ASC ( $r = .60, p = .00; r = .45, p = .03; r = .48, p = .02$  respectively; Figure 4) but there were no significant correlations



between the CD Index and performance in the language measures in typically developed individuals (Figure 5).

We tested whether symptom severity was correlated with the CD index in adults with ASC.

There were no significant correlations between the CD Index and the AQ Likert total score ( $r = .35, p = .11$ ), the ADOS Communication total score ( $r = -.40, p = .09$ ), or the ADOS Social Interaction total score ( $r = -.41, p = .08$ ) but there was a significant correlation between the ADOS Repetitive Interests and Stereotyped behaviours total score ( $r = -.50, p = .03$ ).

(Table 2 here)

(Figure 4 here)

(Figure 5 here)

## Discussion

We examined whether individuals with ASC show generally enhanced discrimination and/or reduced categorical perception of speech stimuli. Individuals with ASC did not show more accurate discrimination at any of the VOT steps in comparison with typically developed adults. This outcome contrasts with a number of previous studies which showed generally enhanced discrimination in basic auditory stimuli (Bonnell et al. 2003; Bonnell et al. 2010; Jones et al. 2009; Foxton et al. 2003; Stewart et al. 2015). It may be that the enhancement shown by individuals with ASC in these studies is specific to particular types of stimuli. The majority of studies have found enhanced discrimination for pitch, either in sinusoidal tones (Bonnell et al. 2003; Jones et al. 2009) or in speech (Heaton 2005; Heaton et al. 2008). One recent study has found that this enhanced discrimination also occurs in the temporal dimension, where

1 participants were required to discriminate between pairs of tones and identify which pair had  
2 the longer gap between them (Stewart et al. 2015). In the current study, we asked whether these  
3 findings extended to VOT, an extension of the temporal dimension, where individuals are  
4 asked to make judgements regarding the temporal lag between the release burst of the  
5 consonant and the onset of the periodic wave of the following vowel. However, no enhanced  
6 discrimination was found at any of the VOT steps. Given that Stewart et al. (2015) used non-  
7 speech stimuli and the current study used speech-like stimuli, it may be that the enhanced  
8 discrimination effect does not apply to the discrimination of temporal acoustic features in the  
9 context of speech perception.  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

24 A second potential explanation for this discrepancy is that previous findings of enhanced  
25 discrimination in individuals with ASC may be a function of enhanced performance on  
26 nonverbal-IQ, as nonverbal-IQ is highly related to stimulus discrimination ability (Deary,  
27 1994). Indeed, enhanced discrimination of auditory materials in ASC was not found in Kargas  
28 et al. (2015), who matched their participants with or without ASC on both age and IQ. The two  
29 groups in our study were also matched person by person on Raven's progressive matrices,  
30 which may account for the lack of general difference in discrimination level between the ASC  
31 and the control group. A strategic decision was taken when designing the study to match the  
32 participants on nonverbal-IQ, and to recruit a sample who were relatively homogeneous for IQ.  
33 This decision means that the findings from this study are somewhat limited in their  
34 generalisability along the dimension of IQ. We suggest that this paradigm is extended to a  
35 group who vary in IQ.  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55

56 We also hypothesised that individuals with ASC would show reduced categorical perception of  
57 phonemes compared to typically developed individuals. In order to assess this, we calculated a  
58  
59  
60  
61  
62  
63  
64  
65

1 categorical discrimination index (CD Index) which assessed whether there was a difference  
2 between perception of phonemes at the edges versus the boundary and we assessed the shape of  
3 the graph. The CD Index was smaller in adults with ASC than in IQ-matched controls - this did  
4 not reach statistical significance in our sample. However, there was a trend, and the effect size  
5 was moderate. Adults with ASC did not have a noticeable peak at the boundary, and their  
6 performance was similar across the VOT steps. The discrimination function was more  
7 platykurtic for adults with ASC than for typically developing individuals. There is variability in  
8 the data with overlap between the groups, but the effect size is moderate, and worthy of  
9 discussion. The pattern of results shown suggests that the perception in individuals with ASC is  
10 less constrained by speech categories than individuals without ASC, a suggestion supported by  
11 Haesen et al. (2011) who propose that individuals with ASC focus on the local, perceptual  
12 features whereas typically developing individuals focus on socially relevant cues such as  
13 speech. The CD Index correlated with identified repetitive behaviours from the ADOS, but not  
14 with other subscales from the ADOS or with autistic traits. The lack of effect must be  
15 interpreted with caution as it may be due to a lack of variability in the ADOS and AQ scores  
16 and the lack of power.

17 One further concern is that the ABX discrimination task taps into attentional, language or  
18 working memory processes and is not assessing discrimination. This seems unlikely, as  
19 participant groups did not differ on reading, spelling, nonword repetition or on lexical  
20 discrimination. The ASC group were also matched one to one on IQ, which reduces the  
21 concern that this group was impaired on general cognitive ability. Crucially, the groups did not  
22 differ in their overall level of discrimination, but only in the shape of the discrimination curve.

Speech perception in individuals without ASC is optimised to differentiate linguistic categories, such as the /k/ sound in ‘coat’ and /g/ sound in ‘goat’, such that the system is relatively insensitive to the acoustic differences that do not contribute to those contrasts. In comparison, speech perception in individuals with ASC is more sensitive to acoustic differences unrelated to linguistic categories. Our findings are consistent with those of DePape et al. (2012). In one of their experiments, DePape and colleagues tested the discrimination of two pairs of foreign language sounds (Zulu consonants) by English speakers: one that maps onto two different native categories and one that maps onto a single native category. Individuals with ASC showed a smaller difference between the two-category and single-category mapping conditions than did controls. The implication is that, much like the participants in our ASC group, the individuals with ASC in DePape et al. (2012) were less influenced by their native speech categories compared to individuals without ASC. There are two main positions regarding phonemic awareness, the first, the accessibility position suggests that phonemic segments are functional and available for basic speech processing tasks from early infancy (Kuhl et al. 1992). The second position suggests that there is more gradual development over the course of childhood (Fowler 1991; Walley 1993). The current study does not test early childhood, nor does it test the relationship between discrimination of speech sounds and language development and therefore does not differentiate between these positions. Rather, the results provide evidence for a reduction in the use of existing categories in speech perception. It remains to be seen whether the effects are due to weaker or delayed category formation in ASC or a less specialised perceptual system in ASC that attends to details that are not relevant to category classification.

We assessed whether the CD Index was related to aspects of language functioning. As the CD Index got larger, performance on vocabulary (Millhill), discrimination of words versus

1 nonwords and reading (WRAT-3 Reading) improved, indicating that the more pronounced the  
2 categorical perception, the better the performance on these tests in individuals with ASC.  
3

4 Between-group comparisons showed that the correlations were significantly larger in the  
5 individuals with ASC than the IQ-matched adults for both vocabulary and reading. The  
6  
7 implication is that some of the language differences found in individuals with ASC may be due  
8  
9 to reduced categorical perception of speech sounds. Our findings suggest the importance of  
10  
11 further examining the link between speech processing and language functioning in ASC. Social  
12  
13 factors, such as, social network size, the demographic make-up of the community and quality  
14  
15 of interaction, influence linguistic skills and perception (Lev-Ari & Peperkamp 2016; Lev-Ari  
16  
17 2017; Ramirez-Esparza et al. 2014). It would be interesting to assess whether there are  
18  
19 particular features of ASC due to the diagnosed differences in social interaction skills which  
20  
21 relate to differences in speech perception.  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33

34 One strength of this study is the homogeneous nature of the sample, and the one to one  
35  
36 matching on IQ between the individuals with ASC and typically developed adults. The effect  
37  
38 size for the CD Index comparison is moderate. To achieve a significant effect where the alpha-  
39  
40 level is set to 5% and power to detect an effect is 80%, a sample of 76 would be required. One  
41  
42 may then argue that our study was underpowered. This study is larger than many studies  
43  
44 assessing perceptual processes in individuals with ASC (e.g. Soulieres et al. 2007; Bonnel et al.  
45  
46 2003; Bonnel et al. 2010; Järvinen-Pasley et al. 2008). This study is the first to assess  
47  
48 categorical speech perception, and suggests that individuals with ASC do not process speech  
49  
50 sounds in the same way as those who are typically developed. In addition, the study identifies  
51  
52 an effect size for this important aspect of perception in ASC.  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

These observations spur investigations into whether children with ASC show similar differences in the categorical perception of speech sounds, at what stage this occurs, and whether this difference in the processing of speech sounds is related to later language development. We know that language skills in school-age children with Autism Spectrum Conditions are predictive of both current function and future outcome (Lord and Paul 1997; Kobayashi et al. 1992; Venter et al. 1992). Yau et al. (2016) suggest that children with ASC have developmental differences in their auditory cortex which influences their ability to process speech and non-speech sounds. If categorical perception of speech turns out to be a reliable predictor of a range of language-related processing abilities, it will provide an important source of information in identifying problems in language functioning at an early stage.

The current study sheds some light on differences in perceptual processing in individuals with ASC. Our findings suggest that it is not simply that individuals with ASC are more perceptually sensitive to the physical differences between stimuli, but that they may show qualitative differences in the way that signals are processed.

### Compliance with Ethical Standards

#### *Funding*

This research was supported by a School of Life Sciences, Heriot-Watt doctoral studentship to one of the authors.

#### *Conflict of interest*

Authors declare that they have no conflict of interest.

*Ethical approval*

All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

*Informed consent*

Informed consent was obtained from all individual participants included in the study.

## References

American Psychiatric Association (2000). *Diagnostic and Statistical Manual of Mental Disorders-Fourth Version-Text Revision (DSM-IV-TR)*. Washington, DC.

Ashwin, C., Baron-Cohen, S., Wheelwright, S., O’Riordan, M., & Bullmore, E. T. (2007). Differential activation of the amygdala and the ‘social brain’ during fearful face-processing in Asperger Syndrome. *Neuropsychologia*, 45(1), 2-14.

Austin, E. J. (2005). Personality correlates of the broader autism phenotype as assessed by the Autism Spectrum Quotient (AQ). *Personality and Individual Differences*, 38, 451-460.

Baron-Cohen, S., Wheelwright, S., Skinner, R., Martin, J., & Clubley, E. (2001). The Autism-Spectrum Quotient (AQ): Evidence from Asperger syndrome/high-functioning autism, males and females, scientists and mathematicians. *Journal of Autism and Developmental Disorders*, 31, 5-17.

Bonnel, A., McAdams, S., Smith, B., Berthiaume, C., Bertone, A., Ciocca, V., et al. (2010). Enhanced pure tone pitch discrimination among persons with autism but not Asperger syndrome. *Neuropsychologia*, 48, 2465-2475.

Bonnel, A., Mottron, L., Peretz, I., Trudel, M., Gallun, E., & Bonnel, A-M. (2003). Enhanced pitch sensitivity in individuals with autism: A signal detection analysis. *Journal of Cognitive Neuroscience*, 15(2), 226-235.

Boraston, Z., Blakemore, S. J., Chilvers, R., & Skuse, D. (2007). Impaired sadness recognition is linked to social interaction deficit in autism. *Neuropsychologia*, 45(7), 1501-1510.

Clark, A., Hughes, P., Grube, M., & Stewart, M. E. (2013). Autistic traits and sensitivity to interference with flavour identification. *Autism Research*, 6, 332-336.

Deary, I. J. (1994). Intelligence and auditory discrimination: Separating processing speed and fidelity of stimulus representation. *Intelligence*, 18, 189-213.

DePape, A-M. R., Hall, G. B. C., Tillmann, B., & Trainor, L. J. (2012). Auditory processing in high-functioning adolescents with autism spectrum disorder. *PLoS One*, 7, e44084.

Fowler, A. E. (1991). How early phonological development might set the stage for phoneme awareness. In S. A. Brady & D. P. Shankweiler (Eds.), *Phonological processes in literacy: A tribute to Isabelle Y. Liberman* (pp. 97-117). Hillsdale, New Jersey: Erlbaum.



1 Foxton, J. M., Stewart, M. E., Barnard, L., Rodgers, J., Young, A. H., O'Brien, G., et al.

2 (2003). Absence of auditory 'global interference' in autism. *Brain*, 126, 2703-2709.

3  
4  
5  
6  
7 Frith, U. (2003). *Autism: Explaining the enigma* (2nd ed.). Oxford, UK: Blackwell Publishers.

8  
9  
10  
11 Fugard, A. J. B., Stewart, M. E., & Stenning, K. (2011). Visual/verbal-analytic reasoning bias  
12 as a function of self-reported autistic-like traits: A study of typically developing individuals  
13 solving Raven's Advanced Progressive Matrices. *Autism*, 15, 327-340.

14  
15  
16  
17  
18  
19  
20  
21 Grube, M., Cooper, F.E., & Griffiths, T.D. (2013). Auditory temporal-regularity processing  
22 correlating with language and literacy skill in early adulthood. *Cognitive Neuroscience*, 4, 3-4.

23  
24  
25  
26  
27  
28  
29 Haesen, B., Boets, B., & Wagemans, J. (2011). A review of behavioural and  
30 electrophysiological studies on auditory processing and speech perception in autism spectrum  
31 disorders. *Research in Autism Spectrum Disorders*, 5(2), 701-714.

32  
33  
34  
35  
36  
37  
38  
39 Hamel, R., & Schmittmann, V. (2006). The 20-minute version as a predictor of the Raven  
40 Advanced Progressive Matrices Test. *Educational and Psychological Measurement*, 66, 1039-  
41 1046.

42  
43  
44  
45  
46  
47  
48  
49 Harnad, S. (1987). *Categorical perception: The groundwork of cognition*. New York:  
50 Cambridge University Press.

51  
52  
53  
54  
55  
56 Heaton, P. (2003). Pitch memory, labelling and disembedding in autism. *Journal of Child*  
57 *Psychology and Psychiatry, and Allied Disciplines*, 44, 543-551.

1  
2 Heaton, P. (2005). Interval and contour processing in autism. *Journal of Autism and*  
3  
4 *Developmental Disorders*, 35, 787-793.  
5  
6  
7

8  
9  
10 Heaton, P., Hermelin, B., & Pring, L. (1998). Autism and pitch processing: a precursor for  
11  
12 savant musical ability? *Music Perception*, 15, 291-305.  
13  
14  
15

16  
17 Heaton, P., Hudry, K., Ludlow, A., & Hill, E. (2008). Superior discrimination of speech pitch  
18  
19 and its relationship to verbal ability in autism spectrum disorders. *Cognitive Neuropsychology*,  
20  
21 25, 771-782.  
22  
23  
24

25  
26 Heaton, P., Pring, L., & Hermelin, B. (1999). A pseudo-savant: A case of exceptional musical  
27  
28 splinter skills. *Neurocase*, 5, 503–509.  
29  
30  
31

32  
33  
34 Hoekstra, R., Bartels, M., Verweij, C. J. H., & Boomsma, D. (2007). Heritability of autistic  
35  
36 traits in the general population. *Archives of Pediatrics and Adolescent Medicine*, 16, 372-377.  
37  
38

39 Järvinen-Pasley, A. M., Pasley, J., & Heaton, P. (2008). Is the linguistic content of speech less  
40  
41 salient than its perceptual features in autism? *Journal of Autism and Developmental Disorders*,  
42  
43 38, 239-248.  
44  
45  
46

47  
48 Jones, C. R. G., Happé, F. G. E., Golden, H., Marsden, A. J. S., Tregay, J., Simonoff, E., et al.  
49  
50 (2009). Reading and arithmetic in adolescents with autism spectrum disorders: Peaks and dips  
51  
52 in attainment. *Neuropsychology*, 23, 718-728.  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65

1 Kargas, N., López, B., Reddy, V., & Morris, P. (2015). The relationship between auditory  
2 processing and restricted repetitive behaviors in adults with autism spectrum disorders. *Journal*  
3 *of Autism and Developmental Disorders*, 45, 658-668.

4  
5  
6  
7  
8  
9 Kjelgaard, M. M., & Tager-Flusberg, H. (2001). An investigation of language impairment in  
10 autism: Implications for genetic subgroups. *Language and Cognitive Processes*, 16, 287-308.

11  
12  
13  
14  
15  
16 Kobayashi, R., Murata, T., & Yoshinaga, K. (1992). A follow-up study of 201 children with  
17 autism in Kyushu and Yamaguchi areas, Japan. *Journal of Autism and Developmental*  
18 *Disorders*, 22, 395-411.

19  
20  
21  
22  
23  
24  
25  
26 Korpilahti, P., Jansson-Verkasalo, E., Mattila, M-L., Kuusikko, S., Suominen, K., Rytty, S., et  
27 al. (2007). Processing of affective speech prosody is impaired in Asperger syndrome. *Journal*  
28 *of Autism and Developmental Disorders*, 37, 1539-1549.

29  
30  
31  
32  
33  
34  
35  
36 Kuhl, P. K., Williams, K. A., Lacerda, F., Stevens, K. N. & Lindblom, B. (1992).  
37 Linguistic experience alters phonetic perception in infants by 6 months of age. *Science*, 255,  
38 606–608.

39  
40  
41  
42  
43  
44  
45  
46 Kujala, T., Lepistö, T., Nieminen-von Wendt, T., Näätänen, P., & Näätänen, R. (2005).  
47 Neurophysiological evidence for cortical discrimination impairment of prosody in Asperger  
48 syndrome. *Neuroscience Letters*, 383(3), 260-265.

1 Lev-Ari, S. (in press). *The influence of social network properties on language processing and*  
 2 *use*. In M. S. Vitevitch (Ed.), *Network Science in Cognitive Psychology*. New York, NY:  
 3  
 4 Routledge.  
 5  
 6  
 7

8  
 9 Lev-Ari, S., & Peperkamp, S. (2016). How the demographic make-up of our community  
 10 influences speech perception. *Journal of the Acoustical Society of America*, 139(6), 3076-3087.  
 11  
 12  
 13  
 14  
 15

16 Liberman, A. M., Cooper, F. S., Shankweiler, D. P., & Studdert-Kennedy, M. (1967).  
 17 Perception of the speech code. *Psychological Review*, 74(6), 431-461.  
 18  
 19  
 20  
 21  
 22  
 23

24 Liberman, A. M., Harris, K. S., Hoffman, H. S., & Griffith, B. C. (1957). The discrimination of  
 25 speech sounds within and across phoneme boundaries. *Journal of Experimental Child*  
 26 *Psychology*, 54, 358-368.  
 27  
 28  
 29  
 30  
 31  
 32  
 33

34 Lord, C., & Paul, R. (1997). *Language and communication in autism*. In D. J. Cohen., & F. R.  
 35 Volkmar (Eds.). *Handbook of autism and pervasive developmental disorders*, (2nd ed). New  
 36 York City, NY: John Wiley and Sons. p. 195-225.  
 37  
 38  
 39  
 40  
 41  
 42

43 Lord, C., Risi, S., Lambrecht, L., Cook, E.H., Leventhal, B.L., DiLavore, P.C., et al. (2000).  
 44 The Autism Diagnostic Observation Schedule–Generic (ADOS Generic): A standard measure  
 45 of social and communication deficits associated with the spectrum of autism. *Journal of Autism*  
 46 *and Developmental Disorders*, 30, 205-223.  
 47  
 48  
 49  
 50  
 51  
 52  
 53  
 54  
 55

56 McCann, J., & Peppé, S. (2003). Prosody in autism spectrum disorders: A critical review.  
 57 *International Journal of Language and Communication Disorders*, 38, 325-350.  
 58  
 59  
 60  
 61  
 62  
 63  
 64  
 65

Meilleur, A-A.S., Berthiaume, C., Bertone, A., & Mottron, L. (2014). Autism-specific covariation in perceptual performances: “g” or “p” factor? *PLoS ONE*, 9, e103781.

Mottron, L., & Burack, J.A. (2001). *Enhanced perceptual functioning in the development of autism*. In J. A. Burack., T. Charman., N. Yirmiya., & P. R. Zelazo (Eds.). *The development of autism perspectives from theory and research*. Mahwah, NJ: Lawrence Erlbaum Associates, p. 131-148.

Mottron, L., Peretz, I., & Menard, E. (2000). Local and global processing of music in high functioning persons with autism: Beyond central coherence? *Journal of Child Psychology and Psychiatry*, 41, 1057-1065.

Plaisted, K. C., O'Riordan, M. A., & Baron-Cohen, S. (1998). Enhanced discrimination of novel, highly similar stimuli by adults with autism during a perceptual learning task. *Journal of Child Psychology and Psychiatry, and Allied Disciplines*, 39, 765-775.

Ramirez-Esparza, N., Garcia-Sierra, A., & Kuhl, P. K. (2014). Look who’s talking: Speech style and social context in language input are linked to concurrent and future speech development. *Developmental Science*, 17, 880-891.

Raven, J. C., Raven, J. E., & Court, J. H. (2003). *Section 4: Advanced progressive matrices. Manual for Raven’s progressive matrices and vocabulary scales*. Oxford, UK: Oxford Psychologists Press.

1 Snowling, M. J. (2000). *Dyslexia*. Blackwell publishing.

2  
3  
4 Soulières, I., Mottron, L., Saumier, D., & Larochelle, S. (2007). Atypical categorical  
5 perception in autism: Autonomy of discrimination? *Journal of Autism and Developmental*  
6  
7  
8  
9  
10 *Disorder*, 37, 481-490.

11  
12  
13  
14 Stewart, M. E., Griffiths, T., & Grube, M. (2015). Autistic traits and enhanced perceptual  
15 representation of pitch and time. *Journal of Autism and Developmental Disorders*. DOI  
16  
17  
18  
19 10.1007/s10803-015-2517-3.  
20  
21

22  
23  
24 Stewart, M. E., & Ota, M. (2008). Lexical effects on speech perception in individuals with  
25  
26  
27 “autistic” traits, *Cognition*, 109, 157-162.  
28

29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000

Walley, A. C. (1993). The role of vocabulary development in children's spoken word recognition and segmentation ability. *Developmental Review, 13*, 286-350.

Wilkinson, G. S. (1993). *Wide Range Achievement Test – Third Revision (WRAT-3)*. Wilmington, DE: The Psychological Corporation.

Woodbury-Smith, M. R., Robinson, J., Wheelwright, S., & Baron-Cohen, S. (2005). Screening adults for Asperger syndrome using the AQ: A preliminary study of its diagnostic validity in clinical practice. *Journal of autism and developmental disorders, 35*(3), 331-335.

Yau, S. H., Brock, J., & McArthur, G. (2016). The relationship between spoken language and speech and nonspeech processing in children with autism: a magnetic event-related field study. *Developmental Science, 19*(5), 834-852.

## Figure Captions

*Figure 1.* Hypothetical discrimination curves for VOT. Line (a): a typical curve with a clear peak around the category boundary. Line (b): a curve with a less clear peak but with a similar general level of sensitivity as (a). Line (c): a curve with a less clear peak and a generally higher level of sensitivity than (a).

*Figure 2.* Discrimination of VOTs by adults with ASC and IQ-matched adults (accuracy). Error bars are standard errors.

*Figure 3.* Discrimination of VOTs by adults with ASC and IQ-matched adults (reaction time). Error bars are standard errors.

*Figure 4.* Scatterplots of the CD Index and (a) WRAT-3 Reading score, (b) Lexical discrimination score, and (c) Millhill vocabulary score for adults with ASC.

*Figure 5.* Scatterplots of the CD Index and (a) WRAT-3 Reading score, (b) Lexical discrimination score, and (c) Millhill vocabulary score for IQ-matched adults.



Figure 1 top




Figure 2 top




Figure 3 top




Figure 4 top




Figure 5 top




Table 1. Descriptive statistics of study measures for adults with ASC and IQ-matched adults

|                                     | Adults with ASC |             |          | IQ-matched adults |             |          |
|-------------------------------------|-----------------|-------------|----------|-------------------|-------------|----------|
|                                     | Mean (SD)       | Range       | <i>n</i> | Mean (SD)         | Range       | <i>n</i> |
| AQ Likert Total***                  | 145.2 (19.9)    | 108 – 184   | 23       | 114.2 (21.9)      | 73 – 144    | 23       |
| Raven's                             |                 |             |          |                   |             |          |
| Raw                                 | 16.6 (5.8)      | 5 – 26      | 23       | 16.6 (5.8)        | 5 – 26      | 23       |
| Standard                            | 30.3 (19.8)     | 1 – 70      | 23       | 31.8 (20.8)       | 1 – 70      | 23       |
| Millhill                            | 19.3 (5.3)      | 10 – 28     | 23       | 20.0 (2.8)        | 13 – 24     | 23       |
| Lexical discrimination (% accurate) | 0.94 (0.03)     | 0.87 – 0.98 | 23       | 0.93 (0.03)       | 0.87 – 0.98 | 23       |
| WRAT-3 Reading                      |                 |             |          |                   |             |          |
| Raw                                 | 35.8 (4.4)      | 28 – 42     | 23       | 35.5 (3.2)        | 29 – 42     | 23       |
| Standard                            | 108.2 (8.7)     | 92 – 120    | 23       | 107.0 (7.5)       | 92 – 120    | 23       |
| WRAT-3 Spelling                     |                 |             |          |                   |             |          |
| Raw                                 | 30.0 (6.9)      | 18 – 39     | 23       | 30.6 (4.3)        | 22 – 36     | 23       |
| Standard                            | 106.8 (14.2)    | 80 – 125    | 23       | 107.6 (9.6)       | 89 – 119    | 23       |
| NWR Average (% accurate)            | 0.90 (0.09)     | 61 – 99     | 19       | 0.90 (0.06)       | 0.79 – 0.98 | 19       |
| NWR Total Correct                   | 8.0 (4.8)       | 0 – 17      | 19       | 7.8 (3.9)         | 3 – 16      | 19       |

Note: AQ=Autism-Spectrum Quotient; WRAT-3=Wide Range Achievement Test – Third Revision; NWR=Nonword Repetition; \*  $p < .001$ .

Table 2. Relationships between the CD Index and the language measures for adults with ASC and IQ-matched adults

|                          | Adults with ASC | IQ-matched adults |
|--------------------------|-----------------|-------------------|
|                          | <i>r</i>        | <i>r</i>          |
| WRAT–3 Reading           | <b>.60**</b>    | -.01              |
| WRAT–3 Spelling          | .38             | -.27              |
| Lexical Discrimination   | <b>.45*</b>     | -.04              |
| Millhill                 | <b>.48*</b>     | -.18              |
| NWR Average (% accurate) | .04             | .21               |
| NWR Total Correct        | .26             | .26               |

Note: WRAT–3=Wide Range Achievement Test – Third Revision; NWR=Nonword Repetition; all correlations, two-tailed; uncorrected for multiple comparisons by language measures; \**p* < .05, \*\**p* < .01.

### Author Note

Mary E. Stewart, Heriot-Watt University, Edinburgh, Scotland, EH14 4AS.

Alexandra M. Petrou, Heriot-Watt University, Edinburgh, Scotland, EH14 4AS.

Mitsuhiko Ota, Philosophy, Psychology and Language Sciences, University of Edinburgh, Edinburgh, Scotland, EH8 9AD.

Alexandra M. Petrou is now at Institute of Neuroscience, Newcastle University, Newcastle Upon Tyne, England, NE1 4LP.

### *Acknowledgements*

We would like to thank first and foremost all the individuals who took part in this study. Andrea Clark for pilot data relating to this study.

### *Funding*

This research was supported by a School of Life Sciences, Heriot-Watt doctoral studentship to Alexandra M. Petrou.

Correspondence concerning this article should be addressed to Dr Alexandra Petrou, Institute of Neuroscience, Newcastle University, Newcastle Upon Tyne, England, NE1 4LP. Email: [alexandra.petrou@ncl.ac.uk](mailto:alexandra.petrou@ncl.ac.uk). Tel: +44 191 282 1380.